	Reg.	No.	••••••	
--	------	-----	--------	--

Name.....

B.Sc. DEGREE (C.B.C.S.S.) EXAMINATION, MARCH 2015

Sixth Semester

Core Course-NUCLEAR AND PARTICLE PHYSICS

[Common for Model I B.Sc. Physics, Model II B.Sc. Physics, B.Sc. Physics, B.Sc. Physics-EEM and B.Sc. Physics-Instrumentation]

Time: Three Hours

Maximum Weight: 25

Part A (Objective Type)

	-		swer an qu ur question	uestions. is carries a weight of 1.			
			Bunch	I			
1.	Isomers are the atoms having:						
	(a)	Same mass number.	(b)	Same atomic number.			
	(c)	Same neutron number.	(d)	Both (a) and (b).			
2,	Which of the following nucleus is more stable?						
	(a)	Oxygen nucleus.	(b)	Xenon nucleus.			
	(c)	Iron nucleus.	(d)	Lead nucleus.			
3. Nuclear force is:							
	(a)	Long range force.	(b)	Electrostatic force.			
	(e)	Short range force.	(d)	All of the above.			
4.	Which	ich of the following decay atomic number remains the same ?					
	(a)	Beta decay.	(b)	Gamma decay.			
	(c)	Alpha decay.	(d)	Both (a) and (b).			
			Bunch	п			
5.	The mi	clear shell model is also name	das:				
	(a)	Electron model.	(b)	Independent particle model.			
	(e)	Atom model.	(d)	None.			
6.	Packing fraction is associated with nucleus in terms of:						
	'(a)	Stability.	(b)	Mass defect.			
	(e)	Nuclear force.	(d)	All of the above.			

 $(4 \times 1 = 4)$

7.	Ionisat	ion chamber is filled with which	of the f	following gas :			
	(a)	Methyl bromide.	(b)	Hydrogen.	E -		
	(c)	Nitrogen.	(d)	None.			
8.	Gamow's theory of alpha decay depends on : **********************************						
	(a)	Classical physics.	(b)	Quantum physics.	100		
	(c)	Both (a) and (b).	(d)	None.			
			Bunch	ш	ail send i i a		
9.	The charge of a neutrino is:						
= 8	(a)	Positive.	(b)	Negative.			
	(c)	Neutral.	(d)	Either positive or negative.			
10.	Internal conversion is a:						
	(a)	Single step process.	(b)	Double step process.			
	(e)	Either single or double.	(d)	Both (a) and (b).			
11.	In a nu	clear reaction which is conserved	i :	robnen granen som			
	(a)	Charge.	(b)	Nucleons.			
	(c)	Parity.	(d)	All of the above.			
12.	Nuclea	r fission in a nuclear reactor is :	M. VE				
	(a)	Chain reaction.	(b)				
	(c)	Uncontrolled chain reaction.	(d)	All of the above.			
			Bunch				
13.	In nucl	ear reactors — neutrons a	re used	A SECULIA SELOTE RECEIVANT POR SELECT			
	(a)	Slow.	(b)	Fast.			
	(c)	Very fast.	(d)	Super fast.			
14.	Which one of the following is a baryon?						
	(a)	Photon.	(b)	Electron.			
	(e)	Sigma.	(d)	Kaon.			
15.	A meson have how many quarks:						
	(a)	One.	(b)	Two.			
	(c)	Three.	(d)	None.	2001		
16.		in value of a baryon is :					
	(a)		(h)	1/2.			
		- 1/2.		None.			
	1,400		of sheet,	The state of the s			

Part B (Short Answer Questions)

Answer any **five** questions. Each question carries a weight of 1.

- 17. Define mass defect.
- 18. Write short note on nuclear density.
- 19. Give few names of nuclear detectors.
- 20. Define mean life of a radioactive element.
- 21. State Geiger-Nuttal law.
- 22. Give one example of artificial radio activity.
- 23. What is a thermonuclear reaction?
- 24. Give the quark model for a proton.

 $(5 \times 1 = 5)$

Part C (Short Essay/Problems)

Answer any four questions. Each question carries a weight of 2.

- 25. Give the meson theory of nuclear force.
- Find the energy release, if two, H² nuclei can fuse together to form ²He₄ nucleus. The binding energy per nucleon of H² and He⁴ is 1.1 MeV and 7.0 MeV respectively.
- 27. Write a note on proton-proton cycle.
- 28. Calculate the Q-value of the following nuclear reaction in MeV :- ${}^{7}N^{14} + 2He^{4} \rightarrow {}_{8}O^{17} + {}_{1}H^{1} + Q$ Where, $N^{14} = 14.003074$, $He^{4} = 4.002604$, $O^{17} = 16.99913$, $H^{1} = 1.007825$.
- 29. Write a note on primary and secondary cosmic rays.
- 30. Describe about radiation hazards and their disposal.

 $(4 \times 2 = 8)$

Part D (Essay)

Answer any two questions. Each question carries a weight of 4.

- 31. Describe the working function of a Bainbridge's mass spectrometer.
- 32. Give the Gamow's theory of alpha decay and explain it in detail.
- 33. Discuss the various types of elementary particles in detail.

 $(2 \times 4 = 8)$