

F 6494

(Pages : 3)

Reg. No.....

Name.....

M.Sc. DEGREE (C.S.S.) EXAMINATION, JANUARY 2015

Third Semester

Faculty of Science

Branch I-(A)—Mathematics

MT 03 C11—MULTIVARIATE CALCULUS AND INTEGRAL TRANSFORMS

(2012 Admission onwards)

Time : Three Hours

Maximum Weight : 30

Part A

*Answer any five questions.
Each question has weight 1.*

1. Write the Fourier series generated by $f \in L([0, p])$ with period p . Also write the formulas for the Fourier co-efficients.
2. Write the inversion formula for Fourier transforms.
3. If $f(x) = \|x\|^2$, then what is the directional derivative $f'(c, u)$ of f at c in the direction of u .
4. Show by an example that a function can have a finite directional derivative $f'(c, u)$ for every u but may fail to be continuous at c .
5. State the inverse function theorem.
6. If $f = u + iv$ is a complex-valued function with a determinant at a point z in C , then show that $J_f(z) = |f'(z)|^2$.
7. Define a k -form in an open set $E \subset R^n$.
8. State Stoke's theorem.

(5 × 1 = 5)

Part B

*Answer any five questions.
Each question has weight 2.*

9. State and prove Weierstrass approximation theorem.
10. Show that $B(p, q) = \frac{\sqrt{p} \sqrt{q}}{\sqrt{(p+q)}}$.

Turn over

11. Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by the equation $f(x, y) = (\sin x \cos y, \sin x \sin y, \cos x \cos y)$. Determine the Jacobian matrix $Df(x, y)$.
12. Prove that if f is differentiable at c , then f is continuous at c .
13. State and prove mean value theorem for differential calculus.
14. Let A be an open subset of \mathbb{R}^n and assume that $f: A \rightarrow \mathbb{R}^n$ has continuous partial derivatives $D_j f_i$ on A . If $J_f(x) \neq 0$ for all x in A , prove that f is an open mapping.
15. For every $f \in \zeta(T^K)$, prove that $L(f) = L'(f)$.
16. Suppose $w = \sum_I b_I(x) dx_I$ is the standard representation of a k -form w in an open set $E \subset \mathbb{R}^n$.

Prove that if $w = 0$ in E , then $b_I^{(x)} = 0$ for every increasing k -index I and for every $x \in E$.

(5 × 2 = 10)

Part C

Answer any three questions.

Each question has weight 5.

17. State and prove the convolution theorem for Fourier transforms.
18. Assume that g is differentiable at a , with total derivative $g'(a)$. Let $b = g(a)$ and assume that f is differentiable at b with total derivative $f'(b)$. Prove that the composition $h = f \circ g$ is differentiable at a and $h'(a) = f'(b) \circ g'(a)$.
19. Let u and v be two real valued functions defined on a subset S of the complex plane. Assume that u and v are differentiable at an interior point c of S and the partial derivatives satisfy the Cauchy-Riemann equations at c . Prove that the function $f = u + iv$ has derivative at c and $f'(c) = D_1 u(c) + i D_1 v(c)$.
20. Prove that if both partial derivatives $D_r f$ and $D_k f$ exists in an n -ball (c, δ) and if both are differential at c , then $D_{r,k} f(c) = D_{k,r} f(c)$.
21. Assume that the second order partial derivatives $D_{i,j} f$ exist in an n -ball $B(a)$ and are continuous at a , where a is a stationary point of f . Let $Q(t) = \frac{1}{2} f''(a, t) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n D_{i,j} f(a) t_i t_j$.

Prove that :

- (a) If $Q(t) > 0$ for all $t \neq 0$, f has a relative minima at a .
- (b) If $Q(t) < 0$ for all $t \neq 0$, f has a relative maxima at a .
- (c) If $Q(t)$ takes both positive and negative values, the f has a saddle point at a .

22. Prove the following :—

- (a) If w and λ are k - and m -forms respectively of class \mathcal{C}^1 in E , then

$$d(w \wedge \lambda) = (dw) \wedge \lambda + (-1)^k w \wedge d\lambda.$$

- (b) If w is of class \mathcal{C}^n in E , then $d^2 w = 0$. Here E is some open set in \mathbb{R}^n .

(3 × 5 = 15)